


09/719, 272



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

B

Search

PubMed

for

Preview

Go

C

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

- Search History will be lost after one hour of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.

Search	Most Recent Queries	Time	Result
#13	Search #11 AND monoclonal antibody	15:58:34	<u>9</u>
#12	Search #11 and monoclonal antibody	15:56:43	<u>196</u>
#11	Related Articles for PubMed (Select 8995282)	15:55:30	<u>196</u>
#10	Search the journal of biological chemistry [jour] AND 272[volume] AND 448[page] Field: Title Word	15:55:26	<u>1</u>

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

Write to the Help Desk

NCBI | NLM | NIH

Department of Health &amp; Human Services

Freedom of Information Act | Disclaimer

Apr 24 2003 12:56:45

[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[PMC](#)[Taxonomy](#)[OMIM](#)[B](#)

Search

PubMed

for

[Limits](#)[Preview/Index](#)[History](#)[Clipboard](#)[Details](#)[About Entrez](#)

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

[Overview](#)[Help | FAQ](#)[Tutorial](#)[New/Noteworthy](#)[E-Utilities](#)

PubMed Services

[Journals Database](#)[MeSH Browser](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[LinkOut](#)[Cubby](#)

Related Resources

[Order Documents](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)[Privacy Policy](#)☐ 1: Br J Cancer 2003 Apr 22;88(8):1223-8[Related Articles, Li](#)**BJC** British Journal of Cancer

## Overexpression of leucocyte common antigen (LAR) P-subunit in thyroid carcinomas.

Konishi N, Tsujikawa K, Yamamoto H, Ishida E, Nakamura M, Shimada K, Yane K, Yamashita H, Noguchi S.

Protein tyrosine phosphatase (PTPase) dephosphorylation and protein tyrosine kinase (PTKs) phosphorylation of key signal transduction proteins may be regulated by extracellular signals, making PTPases important in the regulation of cell proliferation. Leucocyte common antigen (LAR), a receptor-like PTPase, consists of E-subunit, containing the cell adhesion molecule-like receptor region, and P-subunit specific for a short segment of the extracellular region, the transmembrane peptide, and two cytoplasmic PTPase domains. We produced a monoclonal antibody against the LAR P-subunit for immunohistochemical screening of LAR expression in normal and tumorous tissues. Gliomas and gastric, colorectal, lung, breast and prostate cancer showed weak and relatively infrequent expression. Intense and diffuse expression, however, was detected in 95% (227 out of 239) of thyroid carcinomas, but only 12% (22 out of 128) of adenomas and no cases of benign thyroid disease were immunopositive. In contrast to broad staining in carcinomas, LAR expression in thyroid adenomas was often found in small focal or locally invasive areas. Western blot analysis similarly detected LAR P-subunit protein in thyroid carcinomas, but not in normal tissues. We believe this to be the first demonstration of LAR overexpression in thyroid carcinoma and may help to elucidate the role of PTPases in the development of malignancy. British Journal of Cancer (2003) 88, 1223-1228.  
doi:10.1038/sj.bjc.6600876www.bjcancer.com

PMID: 12698188 [PubMed - in process]

## WEST

### Create A Case

Select?	Database	Query	Plural	Op	Thesaurus	Set Name
<input checked="" type="checkbox"/>	USPT	leucocyte common antigen	YES	ADJ	ASSIGNEE	L1
<input checked="" type="checkbox"/>	USPT	LAR	YES	ADJ	ASSIGNEE	L2
<input checked="" type="checkbox"/>	USPT	thyroid carcinoma	YES	ADJ	ASSIGNEE	L3
<input checked="" type="checkbox"/>	USPT	L3 and (L2 or L1)	YES	ADJ	ASSIGNEE	L4
<input checked="" type="checkbox"/>	USPT	(L1 or L2) and antibody	YES	ADJ	ASSIGNEE	L5
<input checked="" type="checkbox"/>	USPT	L5 and carcinoma	YES	ADJ	ASSIGNEE	L6
<input checked="" type="checkbox"/>	USPT	L6 and thyroid	YES	ADJ	ASSIGNEE	L7

Please enter the case name:








### Rules for naming Cases

- Case names can only contain alphanumeric characters including underscore (\_).
- Any other special characters or punctuation characters will be automatically removed prior to saving the case.
- All white space characters will be replaced by an underscore.

09/19, 272

# OMIM

Online Mendelian Inheritance in Man



Johns  
Hopkins  
University

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Search

OMIM

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

Display

Detailed

Show:

20

Send to

Text

**\*179590**

Links

## PROTEIN-TYROSINE PHOSPHATASE, RECEPTOR-TYPE, F; PTPRF

### Alternative titles; symbols

### RECEPTOR-LINKED PROTEIN-TYROSINE PHOSPHATASE LAR LEUKOCYTE ANTIGEN-RELATED TYROSINE PHOSPHATASE; LAR

Gene map locus 1p32

### TEXT

The LAR gene (symbolized PTPRF) encodes a membrane protein that has a cytoplasmic domain with homology to protein-tyrosine phosphatase 1B (176885) and an extracellular domain homologous to the neural cellular adhesion molecule NCAM (116930).

The human LAR molecule closely resembles cell adhesion molecules, which suggests that it may be involved in the regulation of phosphotyrosine levels through cell-cell or cell-matrix interactions. As a first step toward site-directed mutagenesis studies of LAR function, Schaapveld et al. (1995) characterized the mouse *Ptprf* gene. They found that its cytoplasmic region is encoded by 11 exons that span only 4.5 kb of genomic DNA. Compared to the known exon-intron structures of other mammalian receptor-like protein tyrosine phosphatase genes such as *Ptpa* (encoding LRP; 176884) and *Ptpc* (coding for Ly-5; 151460), the portion of the *Ptprf* gene encoding the cytoplasmic region of murine LAR contained not only smaller, but also fewer introns. ⚡

O'Grady et al. (1994) demonstrated that the human LAR gene is composed of 33 exons spanning over 85 kb. Exon 2 encodes the signal sequence and the first 4 amino acids in the mature LAR protein. The 3 immunoglobulin-like domains are encoded by exons 3 to 7, and the 8 fibronectin type III (FN-III) domains by exons 8 to 17. Exons 18 to 22 encode the juxtamembrane and transmembrane domains, and exons 23 to 33 encode the 2 conserved tyrosine phosphatase domains and the entire 3-prime untranslated region. Alternative splicing of LAR mRNA was revealed by RT-PCR analysis. ⚡

Ahmad and Goldstein (1997) reported that LAR is synthesized as an approximately

200-kD protein that is proteolytically processed into 2 noncovalently associated subunits: a 150-kD extracellular (E) subunit that contains the cell adhesion molecule domains, and an 85-kD phosphatase (P) subunit that contains extracellular, transmembrane, and cytoplasmic domains. Ahmad and Goldstein (1997) carried out several studies to elucidate the relationship between LAR and the insulin signaling pathway. They demonstrated that anti-LAR antibodies inhibit the activity of overexpressed human insulin receptor (147670) in Chinese hamster ovary cells. Immunoprecipitation of LAR from cell lysates and immunoblotting with antibody to the insulin receptor (or vice versa) showed a physical association between LAR and the insulin receptor. In insulin-stimulated rat liver cells, LAR was temporally internalized into a similar endosomal compartment as the insulin receptor. Ahmad and Goldstein (1997) concluded that LAR acts as a protein-tyrosine phosphatase that negatively regulates the insulin signaling pathway. ☞

Nam et al. (1999) determined the crystal structure of the tandem D1 and D2 domains of the human LAR.

By in situ hybridization, Disteche et al. (1989) mapped the LAR gene to 1p34-p32. The related leukocyte common antigen (LCA, also known as CD45; 151460) also maps to chromosome 1. By in situ hybridization, Jirik et al. (1992) mapped LAR, a putative tumor suppressor gene, to 1p32, a region frequently deleted in human neuroblastoma and pheochromocytoma. By fluorescence in situ hybridization, Schaapveld et al. (1995) mapped the Ptp<sup>trf</sup> gene to mouse chromosome 4. Harder et al. (1995) found that coamplification of the PTPRF gene and the MYCL1 gene (164850) in a small cell lung cancer line supported close linkage of the 2 genes. ☞

Most receptor-like transmembrane protein-tyrosine phosphatases (PTPases) such as CD45 and the PTPRF molecule have 2 tandemly repeated PTPase domains in the cytoplasmic segment. Tsujikawa et al. (2001) examined the function of each PTPase domain of PTPRF in vivo using a potential physiologic substrate, insulin receptor, and PTPRF mutant proteins. PTPRF associated with and preferentially dephosphorylated the insulin receptor that was tyrosine phosphorylated by insulin stimulation. Its association was mediated by PTPase domain 2, because the cys1813-to-ser mutation in domain 2 resulted in weakening of the association. The cys1522-to-ser mutant protein, which is defective in the PTPRF PTPase domain 1 catalytic site, was tightly associated with tyrosine-phosphorylated insulin receptor, but failed to dephosphorylate it, indicating that PTPRF PTPase domain 1 is critical for dephosphorylation of tyrosine-phosphorylated insulin receptor. The authors concluded that each domain of PTPRF plays distinct functional roles through phosphorylation and dephosphorylation in vivo. ☞

Schaapveld et al. (1997) used gene targeting in mouse embryonic stem cells to generate mice lacking sequences encoding both LAR phosphatase domains. Homozygous mutant mice develop and grow normally. However, mammary glands of homozygous LAR-deficient females were incapable of delivering milk due to an impaired terminal differentiation of alveoli at late pregnancy. The authors concluded that LAR-mediated signaling may play an important role in mammary gland development and function. ☞

## REFERENCES

1. Ahmad, F.; Goldstein, B. J. :  
Functional association between the insulin receptor and the transmembrane protein-tyrosine phosphatase LAR in intact cells. *J. Biol. Chem.* 272: 448-457, 1997.  
PubMed ID : [8995282](#)
2. Disteche, C. M.; Adler, D. A.; Tedder, T. F.; Saito, H. :  
Mapping of the genes for LYAM1, a new lymphocyte adhesion molecule, and for LAR, a new receptor-linked protein tyrosine phosphatase, to human chromosome 1 (Abstract) *Cytogenet. Cell Genet.* 51: 990 only, 1989.
3. Harder, K. W.; Saw, J.; Miki, N.; Jirik, F. :  
Coexisting amplifications of the chromosome 1p32 genes (PTPRF and MYCL1) encoding protein tyrosine phosphatase LAR and L-myc in a small cell lung cancer line. *Genomics* 27: 552-553, 1995.  
PubMed ID : [7558042](#)
4. Jirik, F. R.; Harder, K. W.; Melhado, I. G.; Anderson, L. L.; Duncan, A. M. V. :  
The gene for leukocyte antigen-related tyrosine phosphatase (LAR) is localized to human chromosome 1p32, a region frequently deleted in tumors of neuroectodermal origin. *Cytogenet. Cell Genet.* 61: 266-268, 1992.  
PubMed ID : [1486801](#)
5. Nam, H.-J.; Poy, F.; Krueger, N. X.; Saito, H.; Frederick, C. A. :  
Crystal structure of the tandem phosphatase domains of RPTP LAR. *Cell* 97: 449-457, 1999.  
PubMed ID : [10338209](#)
6. O'Grady, P.; Krueger, N. X.; Streuli, M.; Saito, H. :  
Genomic organization of the human LAR protein tyrosine phosphatase gene and alternative splicing in the extracellular fibronectin type-III domains. *J. Biol. Chem.* 269: 25193-25199, 1994.  
PubMed ID : [7929208](#)
7. Schaapveld, R. Q.; Schepens, J. T.; Robinson, G. W.; Attema, J.; Oerlemans, F. T.; Fransen, J. A.; Streuli, M.; Wieringa, B.; Hennighausen, L.; Hendriks, W. J. :  
Impaired mammary gland development and function in mice lacking LAR receptor-like tyrosine phosphatase activity. *Dev. Biol.* 188: 134-146, 1997.  
PubMed ID : [9245518](#)
8. Schaapveld, R. Q. J.; van den Maagdenberg, A. M. J. M.; Schepens, J. T. G.; Olde Weghuis, D.; Geurts van Kessel, A.; Wieringa, B.; Hendriks, W. J. A. J. :  
The mouse gene *Ptpfr* encoding the leukocyte common antigen-related molecule LAR: cloning, characterization, and chromosomal localization. *Genomics* 27: 124-130, 1995.  
PubMed ID : [7665159](#)

9. Tsujikawa, K.; Kawakami, N.; Uchino, Y.; Ichijo, T.; Furukawa, T.; Saito, H.; Yamamoto, H. :

Distinct functions of the two protein tyrosine phosphatase domains of LAR (leukocyte common antigen-related) on tyrosine dephosphorylation of insulin receptor. *Molec. Endocr.* 15: 271-280, 2001.

PubMed ID : [11158333](#)

## CONTRIBUTORS

John A. Phillips, III – updated : 7/31/2001

Stylianios E. Antonarakis – updated : 7/2/1999

Rebekah S. Rasooly – updated : 10/7/1998

Rebekah S. Rasooly – updated : 4/9/1998

## CREATION DATE

Victor A. McKusick : 10/19/1988

## EDIT HISTORY

cwells : 8/1/2001

cwells : 7/31/2001

mgross : 7/9/1999

kayiaros : 7/2/1999

alopez : 10/23/1998

alopez : 10/23/1998

alopez : 10/7/1998

dkim : 7/23/1998

psherman : 4/10/1998

alopez : 4/9/1998

mark : 12/4/1995

mark : 7/30/1995

terry : 7/24/1995

carol : 1/4/1995

carol : 4/14/1993

supermim : 3/16/1992

carol : 2/21/1992

Copyright © 1966–2003 Johns Hopkins University

<b>Display</b>	Detailed	Show:	20	<b>Send to</b>	Text
----------------	----------	-------	----	----------------	------

[Disclaimer](#) | [Write to the Help Desk](#) | [Privacy Policy](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

09/7/9, 272

7/9/1 (Item 1 from file: 5)  
DIALOG(R) File 5: Biosis Previews(R)  
(c) 2003 BIOSIS. All rts. reserv.

09536218 BIOSIS NO.: 199497544588

**Mutational analysis of proprotein processing, subunit association, and shedding of the LAR transmembrane protein tyrosine phosphatase.**

AUTHOR: Serra-Pages Carles(a); Saito Haruo; Streuli Michel

AUTHOR ADDRESS: (a) Div. Tumor Immunol., Dana Farber Cancer Inst., Boston, MA 02115\*\*USA

JOURNAL: Journal of Biological Chemistry 269 (38):p23632-23641 1994

ISSN: 0021-9258

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

**ABSTRACT:** The **LAR** transmembrane protein tyrosine phosphatase (PTPase) is expressed on the cell surface as a complex of two noncovalently associated subunits derived from a proprotein. The 150-kDa E-subunit contains most of the extracellular region, including the immunoglobulin-like and fibronectin type-III-like domains, whereas the 85-kDa **P - subunit** contains a short ectodomain, the transmembrane peptide, and the two intracellular PTPase domains. The **LAR** extracellular region is released from the cell surface, suggesting that shedding may be a mechanism to regulate IAR PTPase function. Functional regions necessary for IAR proprotein processing, subunit association, and shedding were determined by analyzing the effect of amino acid substitutions of residues surrounding the cleavage site and scanning the **P - subunit** ectodomain. Three amino acid residues were identified, two within a penta-arginine sequence and one C-terminal to the cleavage site, that are essential for efficient **LAR** proprotein cleavage. Several noncontiguous amino acid residues were also identified that play an essential role in **LAR** subunit association. **LAR** shedding is shown to be a consequence of proteolytic cleavage at a second site within the **P - subunit** ectodomain near the transmembrane peptide.

Set	Items	Description
S1	2703	LAR
S2	0	TYROSINE PHOPPHATASE
S3	560	TYROSINE PHOSPHATASE
S4	72	P(W) SUBUNIT
S5	0	S1 AND S3 AND S4
S6	5	S1 AND S4
S7	3	RD (unique items)



1/9/1 (Item 1 from file: 5)  
DIALOG(R) File 5: Biosis Previews(R)  
(c) 2003 BIOSIS. All rts. reserv.

09096593 BIOSIS NO.: 199497104963

Characterization of AE-6 monoclonal antibody recognizing VHCSAGV sequence  
in CD45 PTPase domain.

AUTHOR: Takeuchi Tsutomu; Sekine Hiromi; Koide Jun; Abe Tohru

AUTHOR ADDRESS: Saitama Med. Sch., Saitama\*\*Japan

JOURNAL: Tissue Antigens 42 (4):p441 1993

CONFERENCE/MEETING: 5th International Conference on Human Leukocyte

Differentiation Antigens Boston, Massachusetts, USA November 3-7, 1993

ISSN: 0001-2815

RECORD TYPE: Citation

LANGUAGE: English

REGISTRY NUMBERS: 79747-53-8: PROTEIN TYROSINE PHOSPHATASE

DESCRIPTORS:

MAJOR CONCEPTS: Biochemistry and Molecular Biophysics; Blood and  
Lymphatics (Transport and Circulation); Cell Biology; Clinical  
Immunology (Human Medicine, Medical Sciences); Enzymology (Biochemistry  
and Molecular Biophysics); Membranes (Cell Biology)

BIOSYSTEMATIC NAMES: Hominidae--Primates, Mammalia, Vertebrata, Chordata,  
Animalia

ORGANISMS: human (Hominidae)

BIOSYSTEMATIC CLASSIFICATION (SUPER TAXA): animals; chordates; humans;  
mammals; primates; vertebrates

CHEMICALS & BIOCHEMICALS: PROTEIN TYROSINE PHOSPHATASE

MISCELLANEOUS TERMS: FIBRONECTIN DOMAIN; MEETING ABSTRACT; PROTEIN  
TYROSINE PHOSPHATASE; SIGNAL TRANSDUCTION; TISSUE ANTIGEN

CONCEPT CODES:

02508 Cytology and Cytochemistry-Human

10064 Biochemical Studies-Proteins, Peptides and Amino Acids

10506 Biophysics-Molecular Properties and Macromolecules

10508 Biophysics-Membrane Phenomena

10806 Enzymes-Chemical and Physical

10808 Enzymes-Physiological Studies

15008 Blood, Blood-Forming Organs and Body Fluids-Lymphatic Tissue and  
Reticuloendothelial System

34508 Immunology and Immunochemistry-Immunopathology, Tissue Immunology

00520 General Biology-Symposia, Transactions and Proceedings of  
Conferences, Congresses, Review Annuals

10068 Biochemical Studies-Carbohydrates

BIOSYSTEMATIC CODES:

86215 Hominidae